

Thursday 13 June 2013 – Afternoon

A2 GCE PHYSICS B (ADVANCING PHYSICS)

G495/01 Field and Particle Pictures

Candidates answer on the Question Paper.

OCR supplied materials:

- Data, Formulae and Relationships Booklet (sent with general stationery)
- Insert (inserted)

Other materials required:

- Electronic calculator
- Ruler (cm/mm)

Duration: 2 hours



Candidate forename		Candidate surname	
--------------------	--	-------------------	--

Centre number						Candidate number				
---------------	--	--	--	--	--	------------------	--	--	--	--

INSTRUCTIONS TO CANDIDATES

- The Insert will be found in the centre of this document.
- Write your name, centre number and candidate number in the boxes above. Please write clearly and in capital letters.
- Use black ink. HB pencil may be used for graphs and diagrams only.
- Answer **all** the questions.
- Read each question carefully. Make sure you know what you have to do before starting your answer.
- Write your answer to each question in the space provided. Additional paper may be used if necessary but you must clearly show your candidate number, centre number and question number(s).
- Do **not** write in the bar codes.

INFORMATION FOR CANDIDATES

- The number of marks is given in brackets [] at the end of each question or part question.
- The total number of marks for this paper is **100**.
- You may use an electronic calculator.
-  Where you see this icon you will be awarded marks for the quality of written communication in your answer.
This means for example, you should:
 - ensure that text is legible and that spelling, punctuation and grammar are accurate so that the meaning is clear
 - organise information clearly and coherently, using specialist vocabulary when appropriate.
- You are advised to show all the steps in any calculations.
- This document consists of **24** pages. Any blank pages are indicated.
- The questions in Section C are based on the material in the Insert.

Answer **all** the questions.

SECTION A

1 Here is a list of units:

$$\text{Js}^{-1}$$

$$\text{Jkg}^{-1}$$

$$\text{Jm}^{-1}$$

$$\text{JC}^{-1}$$

State the equivalent unit from the list for

(a) Sv, radiation dose equivalent

..... [1]

(b) Wbs^{-1} , rate of change of flux linkage.

..... [1]

2 The charge on a u quark is $+\frac{2}{3}e$.

The charge on a d quark is $-\frac{1}{3}e$.

Draw lines from the descriptions given in the left hand boxes to the names of the particles given in the right hand boxes.

particle composed of 2 u and 1 d quark

neutron

negatively charged lepton

electron

neutral lepton

proton

neutral hadron

neutrino

[2]

- 3 The rest energy of an electron is about 510 keV.

An electron is accelerated through a potential difference of 90 keV.

Choose the value from the list below which gives the best estimate for the relativistic factor γ of the accelerated electron.

0.80

1.0


1.2

1.4

best estimate = [1]

- 4 Fig. 4.1 shows a length of wire carrying a current at right angles to a magnetic field.

wire of length 0.15 m
carrying current of 0.41 A



uniform magnetic field
of 0.12 T

Fig. 4.1

Calculate the force on the wire.

force = N [2]

- 5 Fig. 5.1 shows the variation of electric field strength E with distance r from a point charge.




Fig. 5.1

Use the graph to estimate the potential difference between $r = 0.3\text{m}$ and $r = 0.7\text{m}$. Describe the method you use.

potential difference = V [3]

6 Fig. 6.1 shows the path of a proton as it is deflected by a nucleus. This is labelled proton **A**.




Fig. 6.1

- (a) Draw an arrow to show the direction of the force acting on the proton at point **Y**. [1]
- (b) Proton **B** has the same energy as proton **A**. Complete the line showing the path of proton **B**. [2]

7 Here is a list of numbers.

- 8 10 12 14 16

Choose from the list the value that gives the best estimate for

- (a) the average emf generated when the magnetic flux through a 400 turn coil changes by 2.5×10^{-4} Wb in 0.012 s

best estimate = V [1]

- (b) the pd across the secondary coil of a transformer when the primary coil is connected to a 230V, 50Hz supply.

number of turns on primary coil = 1200

number of turns on secondary coil = 60

best estimate = V [1]

- 8 A simple model of an atom represents an electron as a standing wave in a box. The wave for the lowest energy level of the electron ($n = 1$) is shown in Fig. 8.1.




Fig. 8.1

The wavelength λ of the electron is given by

$$\lambda = \frac{h}{p}$$

where p is the momentum of the electron.

Momentum is related to kinetic energy E by the equation

$$E = \frac{p^2}{2m}.$$

- (a) Show that $E = \frac{h^2}{2m\lambda^2}$.

[1]

- (b) The wave for the second energy level of the electron ($n = 2$) is shown in Fig. 8.2.




Fig. 8.2

Choose the value from the list below which gives the ratio $\frac{\text{energy for } n = 2}{\text{energy for } n = 1}$ using this model.

0.25 0.5 1.0 2.0 4.0

ratio = [1]

- 9 Fig. 9.1 shows some equipotential lines in a region of electric field.




Fig. 9.1

- (a) Explain how the diagram shows that the electric field is weaker at **A** than at **B**.

[2]

- (b) Draw an arrow through point **X** to show the direction of the field at that point.

[1]

BLANK PAGE

PLEASE DO NOT WRITE ON THIS PAGE

SECTION B

- 10 This question is about an electric motor which uses a rotating magnetic field.

Fig. 10.1 shows the magnetic flux in the motor in which a permanent magnet is the rotor. The flux is shown at one particular instant.




Fig. 10.1

- (a) (i) State how the position of the poles shown in Fig. 10.1 indicate that the permanent magnet will turn clockwise.

[1]

- (ii) State how the shape of the flux lines shown in Fig. 10.1 **also** indicates that the permanent magnet will turn clockwise.

[1]

- (b) The force on the permanent magnet can be increased by increasing the permeance of the system. Suggest **two** ways in which the permeance of the system can be increased.

[2]

- (c) The flux in the stator is produced by 25 Hz alternating currents in the two pairs of coils **X** and **Y**. Fig. 10.2 shows the flux in the stator at times $t = 0.0\text{ s}$ and $t = 10\text{ ms}$.




Fig. 10.2

Fig. 10.3 shows how the alternating current in the **X** and **Y** coils varies with time.




Fig. 10.3

(i) Use Fig. 10.3 to explain the flux change shown in Fig. 10.2.

[2]

(ii) Explain how, as the alternating currents vary over one complete cycle, the flux rotates by one turn.

[2]

(d) When the permanent magnet is replaced with a non-magnetic copper rotor the motor still spins as the flux rotates.

By considering the changing magnetic field in the rotor, explain why the copper rotor spins.

[3]

[Total: 11]

- 11 This question is about a nuclear fusion reaction that may be able to provide a future energy source. The symbol equation for the first stage of the reaction is given below:



The word equation for the reaction is:

deuterium nucleus + tritium nucleus \rightarrow helium nucleus + neutron.

- (a) (i) Here are some data for the reaction:

mass of deuterium nucleus = 2.0135 u

mass of tritium nucleus = 3.0155 u

mass of helium nucleus = 4.0015 u

mass of neutron = 1.0087 u

Show that the total mass of the particles decreases by about 3×10^{-29} kg in each fusion reaction.

$$1 \text{ u} = 1.66 \times 10^{-27} \text{ kg}$$

[2]

- (ii) Calculate the energy released in one reaction.

$$c = 3.0 \times 10^8 \text{ ms}^{-1}$$

energy released = J [2]

- (b) The separation of the deuterium and tritium nuclei must be about 10^{-14} m for fusion to occur.

Show that the work done in bringing the two nuclei to within 10^{-14} m is about 2×10^{-14} J.

$$\text{electric force constant } \frac{1}{4\pi\epsilon_0} = 9.0 \times 10^9 \text{ Nm}^2\text{C}^{-2}$$

$$e = 1.6 \times 10^{-19} \text{ C}$$

[2]

- (c) In a working fusion reactor, a tritium-deuterium plasma is kept at high temperatures to give the nuclei the energy required for fusion to occur.

- (i) Use $E \approx kT$ to estimate the temperature required of the tritium and deuterium plasma for fusion to occur.

$$\text{minimum energy of nuclei required for fusion} = 1 \times 10^{-14} \text{ J}$$

$$\text{Boltzmann constant } k = 1.4 \times 10^{-23} \text{ JK}^{-1}$$


temperature = K [1]

- (ii) Explain why this fusion reaction may occur when the plasma is at a lower temperature.

[2]

[Total: 9]

- 12 This question is about technetium-99m, an isotope that is used in about 20 million medical investigations each year. The isotope is represented as ${}_{43}^{99m}\text{Tc}$. The nucleus falls to a lower energy state with the release of a gamma photon:



The half life of this process is 6 hours.

- (a) Show that λ , the decay constant for the process, is about $3 \times 10^{-5} \text{ s}^{-1}$.

[2]

- (b) In a medical investigation a patient is injected with a compound containing technetium-99m. The original activity of the injected compound is $9.3 \times 10^8 \text{ Bq}$. Each gamma photon released in the decay process has an energy of 140 keV.

Calculate the initial number of technetium-99m nuclei present in the dose and use this value to show that the total energy emitted by the technetium-99m is less than one joule.

$$1 \text{ eV} = 1.6 \times 10^{-19} \text{ J}$$

original number of technetium-99m nuclei =

total energy of dose = J
[4]

- (c) About 99% of the gamma photons produced pass out through the body of the patient. This can yield information of medical value. However, the patient still receives a significant dose.

Use your answer to (b) to calculate an estimate for the whole-body dose equivalent in Sv received by a patient undergoing a technetium-99m investigation.

mass of patient = 77 kg
quality factor of γ radiation = 1

dose equivalent = Sv [3]

- (d) The risk of developing cancer from radiation exposure is approximately 5% per sievert. Use your answer from (c) to consider the risk implications of this procedure for an individual patient and the global effect of the 20 million procedures each year which use technetium-99m.



Make each step in your calculation and explanation clear.

[4]

[Total: 13]

- 13 This question is about the principles behind a mass spectrometer, an instrument which can identify isotopes from very small samples of their ions. Fig. 13.1 shows the basic components of the instrument.




Fig. 13.1

- (a) The velocity selector contains two plates separated by 0.022 m. There is a potential difference of 440V between the plates, producing a uniform electric field.

- (i) Show that a singly-charged positive ion will experience a force of 3.2×10^{-15} N due to the electric field between the plates.

$$\text{charge on ion} = 1.6 \times 10^{-19} \text{ C}$$

[1]

- (ii) A uniform magnetic field of 0.11 T also acts between the plates as shown in the diagram.

The force F on a charge q moving at speed v in a magnetic field of strength B is given by $F = qvB$. With the aid of a suitable calculation, explain why the path of a singly-charged positive ion moving at about $1.8 \times 10^5 \text{ ms}^{-1}$ can be a straight line through the velocity selector.

[3]

(b) Singly-charged tin ($^{118}\text{Sn}^+$) ions enter the magnetic deflection region as shown in Fig. 13.2.




Fig. 13.2

(i) The magnetic force on the ions acts centripetally.

Show that the radius of the path of an ion of charge q and mass m moving at velocity v in a magnetic field of strength B is given by $r = \frac{mv}{Bq}$.

[1]

(ii) Calculate the radius of the path followed by a singly-charged $^{118}\text{Sn}^+$ ion of mass $2.0 \times 10^{-25} \text{ kg}$ with a speed of $1.8 \times 10^5 \text{ ms}^{-1}$. The magnetic field in the deflection region is 0.70 T .

radius = m [2]

(iii) The sample of tin also includes more massive singly-charged $^{120}\text{Sn}^+$ ions which also pass through the velocity selector at a speed of $1.8 \times 10^5 \text{ ms}^{-1}$. Describe and explain how their path through the magnetic deflection region differs from the path of $^{118}\text{Sn}^+$ ions.

[2]

[Total: 9]

18
SECTION C

These questions are based on the Insert.

- 14 Fig. 14.1 shows the fundamental standing wave pattern of a length of steel violin string.




Fig. 14.1

- (a) The length, L , of the string is 0.32 m. State the wavelength of the fundamental standing wave shown.

wavelength = m [1]

- (b) The frequency of the fundamental note produced is 440 Hz. Calculate the speed of the wave along the string.

speed = ms^{-1} [2]

- (c) The speed v of a transverse wave along a stretched string like this is given by the equation

$$v = \sqrt{\frac{T}{\mu}}$$

where T is the tension and μ the mass per unit length of the string.

Show that the units of $\sqrt{\frac{T}{\mu}}$ are ms^{-1} .

- (d) (i) Show that the frequency of the fundamental mode of oscillation is given by

$$f = \frac{1}{2L} \sqrt{\frac{T}{\mu}}$$

where L is the length of the string.

[1]

- (ii) Show that if the tension T were increased by 10% (a factor of 1.1) then the frequency of the fundamental note would be about 460 Hz. Assume that both μ and L remain constant.

[2]

- (iii) In practice, the value of μ will change as the tension is increased. Explain why.

[2]

- (iv) Suggest and explain how the change in μ affects the value of the frequency you have calculated in part (ii).

[2]

[Total: 12]

15 In line 30 in the article it is stated that the air inside the instrument can resonate.

(a) Using a suitable example, explain the meaning of *resonance*.

[2]

(b) The vibrations lead to standing waves in the body of the instrument (Chladni patterns, line 36 in the article).

Explain how the sand displays these standing wave patterns.

[2]

(c) The material from which a traditional (acoustic) violin is made significantly affects the sound it makes. Explain how making the violin body from stiffer wood would affect the resonant behaviour of the instrument.

[2]

[Total: 6]

16 A diagram of a magnetic pickup for a violin string is shown in Fig. 16.1.




Fig. 16.1

(a) Add **two** lines of flux to the figure to indicate the magnetic circuit created by the pickup. [2]

(b) The permanent magnet used is 6.0 mm in diameter.

Show that if the field strength at the poles of the magnet is 0.10 T, the magnetic flux in those regions is about 3×10^{-6} Wb.

[2]

(c) The coil of wire consists of 8200 turns. Show that this requires about 150 m of wire.

[2]

- (d) The wire has a resistance of 50Ω per metre. An emf of 0.12V is induced across the coil. Calculate the current in the coil when the ends of the wire are joined together.

current = A [2]

- (e) Explain how a steel wire vibrating near the pole of the magnet causes an emf to be induced in the coil.



Your explanation should be clear and follow a logical order.

[4]

[Total: 12]

- 17 (a) Explain how a piezoelectric pickup detects the vibrations of the strings and explain why the strings do not have to be steel when such a pickup is used.

[4]

- (b) Suggest an advantage of using a piezoelectric pickup rather than a magnetic pickup for a violin.

[1]

[Total: 5]

Question 18 begins on page 24

18 Fig. 18.1 shows the waveform of a pure note. Fig. 18.2 shows a more complex note.




Fig. 18.1




Fig. 18.2

(a) Use Fig. 18.1 to calculate the frequency of the pure note.

frequency = Hz [1]

(b) Explain why both notes have the same pitch yet sound different.

[2]

[Total: 3]

END OF QUESTION PAPER



Copyright Information

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact the Copyright Team, First Floor, 9 Hills Road, Cambridge CB2 1GE.

OCR is part of the Cambridge Assessment Group; Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.