

**Tuesday 26 June 2018 – Morning**

**A2 GCE MATHEMATICS**

**4734/01 Probability & Statistics 3**

**QUESTION PAPER**



Candidates answer on the Printed Answer Book.

**OCR supplied materials:**

- Printed Answer Book 4734/01
- List of Formulae (MF1)

**Duration: 1 hour 30 minutes**

**Other materials required:**

- Scientific or graphical calculator

**INSTRUCTIONS TO CANDIDATES**

These instructions are the same on the Printed Answer Book and the Question Paper.

- The Question Paper will be found inside the Printed Answer Book.
- Write your name, centre number and candidate number in the spaces provided on the Printed Answer Book. Please write clearly and in capital letters.
- **Write your answer to each question in the space provided in the Printed Answer Book.** If additional space is required, you should use the lined page(s) at the end of the Printed Answer Book. The question number(s) must be clearly shown.
- Use black ink. HB pencil may be used for graphs and diagrams only.
- Answer **all** the questions.
- Read each question carefully. Make sure you know what you have to do before starting your answer.
- Do **not** write in the barcodes.
- You are permitted to use a scientific or graphical calculator in this paper.
- Give non-exact numerical answers correct to 3 significant figures unless a different degree of accuracy is specified in the question or is clearly appropriate.

**INFORMATION FOR CANDIDATES**

This information is the same on the Printed Answer Book and the Question Paper.

- The number of marks is given in brackets [ ] at the end of each question or part question on the Question Paper.
- **You are reminded of the need for clear presentation in your answers.**
- The total number of marks for this paper is **72**.
- The Printed Answer Book consists of **12** pages. The Question Paper consists of **4** pages. Any blank pages are indicated.

**INSTRUCTION TO EXAMS OFFICER/INVIGILATOR**

- Do not send this Question Paper for marking; it should be retained in the centre or recycled. Please contact OCR Copyright should you wish to re-use this document.

Answer **all** the questions.

- 1 The random variables  $X$  and  $Y$  have independent Poisson distributions with parameters 2 and 3 respectively, and  $Z = 3X + 4Y$ . Find  $P(Z = E(Z))$ . [5]
- 2 A 95% confidence interval for the mean  $\mu$  of a certain population, based on a sample of size 35, is [6.0061, 7.9939]. Find the minimum sample size such that the width of a 95% confidence interval for  $\mu$  is less than 1. [4]
- 3 The cumulative density function of the continuous random variable  $X$  is given by

$$F(x) = \begin{cases} 0 & x < 1, \\ \frac{2x-2}{x+3} & 1 \leq x \leq 5, \\ 1 & x > 5. \end{cases}$$

Given that  $Y = 2X - 3$ , find the probability density function of  $Y$ . [6]

- 4 The mean number of matches in a box is claimed to be 48. A random sample of 7 boxes is taken and the number of matches in each box is counted. The results are shown below.

48 45 46 47 47 46 48

Stating a necessary assumption, test at the 2.5% significance level whether the mean number of matches in a box is less than 48. [8]

- 5 A certain brand of ice cream is sold in cartons of different sizes. Large cartons contain ice cream whose mass is normally distributed with mean 412 g and standard deviation 10 g. Small cartons contain ice cream whose mass is normally distributed with mean 112 g and standard deviation 8 g.
- (i) Find the probability that the total mass of ice cream in two randomly chosen large cartons and two randomly chosen small cartons is greater than 1 kg. [4]
- (ii) Find the probability that the mass of ice cream in a randomly chosen large carton is greater than 4 times the mass of ice cream in a randomly chosen small carton. [4]

- 6 The continuous random variable  $X$  has probability density function

$$f(x) = \begin{cases} k \cos x & 0 \leq x < \frac{\pi}{4}, \\ k \sin x & \frac{\pi}{4} \leq x \leq \frac{\pi}{2}, \\ 0 & \text{otherwise.} \end{cases}$$

- (i) Show that  $k = \frac{1}{\sqrt{2}}$ . [3]
- (ii) Find  $P(X \leq 1)$ . [3]
- (iii) Find the upper quartile of  $X$ . [3]

- 7 Greyhound racing in England involves six dogs racing over distances of approximately 500 m. The dogs wear jackets numbered from 1 to 6. A researcher observes that the winning dog in exactly 24 of 80 randomly chosen races wore jacket number 3.

(i) The probability that a randomly chosen race is won by the dog wearing jacket number 3 is denoted by  $p$ . Calculate an approximate 99% confidence interval for  $p$ . [3]

(ii) Explain whether your result from part (i) is consistent with  $p$  taking the value 0.2. [1]

Greyhound racing meetings in England usually consist of 12 races. The researcher chooses 100 such meetings at random and, at each meeting, he records how many of the 12 races were won by the dog wearing jacket number 3. The results for the 100 meetings are given in the table below.

Number of races won by the dog wearing jacket number 3	0	1	2	3	4	5	6	7	8 or more
Observed frequency	9	18	33	17	14	6	2	1	0

(iii) Show that the proportion of races won by the dog wearing jacket number 3 at the 100 meetings was 0.2. [2]

The expected frequencies using a binomial distribution with  $n = 12$  are given in the table below.

Number of races won by the dog wearing jacket number 3	0	1	2	3	4	5	6	7	8 or more
Expected frequency	6.87	20.62	28.35	23.62	13.29	5.32	1.55	0.33	0.05

(iv) Show how for the case where 4 races were won by dog number 3 the expected frequency of 13.29 was calculated. [2]

(v) Carry out a  $\chi^2$  test, at the 5% significance level, to test the null hypothesis that the data can be well modelled by a binomial distribution with  $n = 12$ . [6]

- 8 The numbers of hours of sunshine observed on 8 randomly chosen days in London were as follows.

9.3 3.9 11.8 5.0 10.6 0.0 6.1 2.7

The numbers of hours of sunshine observed on 6 randomly chosen days in Berlin were as follows.

7.3 2.5 6.0 9.8 12.1 4.6

It may be assumed that the number of hours of sunshine in each city is normally distributed.

(i) Carry out a suitable  $t$ -test, at the 1% significance level, to test whether the mean daily number of hours of sunshine in Berlin is higher than the mean daily number of hours of sunshine in London. [9]

(ii) Explain how the observations could be modified so as to use a paired-sample  $t$ -test, and explain why this would be preferable. [2]

- 9 The results of an examination are represented in the contingency table below, in which  $x$  and  $y$  are integers.

	Pass	Fail	Total
Females	$x$	$y$	50
Males	$y$	$x$	50
Totals	50	50	100

The null hypothesis, that there is no association between examination result and gender, was rejected at the 5% significance level. Given that  $x > y$ , find the smallest possible value of  $x$ . [7]

### END OF QUESTION PAPER



Oxford Cambridge and RSA

#### Copyright Information

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website ([www.ocr.org.uk](http://www.ocr.org.uk)) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact the Copyright Team, First Floor, 9 Hills Road, Cambridge CB2 1GE.

OCR is part of the Cambridge Assessment Group; Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.