

GCE

Further Mathematics B MEI

Y434/01: Numerical methods

A Level

Mark Scheme for June 2023

OCR (Oxford Cambridge and RSA) is a leading UK awarding body, providing a wide range of qualifications to meet the needs of candidates of all ages and abilities. OCR qualifications include AS/A Levels, Diplomas, GCSEs, Cambridge Nationals, Cambridge Technicals, Functional Skills, Key Skills, Entry Level qualifications, NVQs and vocational qualifications in areas such as IT, business, languages, teaching/training, administration and secretarial skills.

It is also responsible for developing new specifications to meet national requirements and the needs of students and teachers. OCR is a not-for-profit organisation; any surplus made is invested back into the establishment to help towards the development of qualifications and support, which keep pace with the changing needs of today's society.

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by examiners. It does not indicate the details of the discussions which took place at an examiners' meeting before marking commenced.

All examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the report on the examination.

© OCR 2023

Text Instructions

1. Annotations and abbreviations

Annotation in scoris	Meaning
√and ×	
BOD	Benefit of doubt
FT	Follow through
ISW	Ignore subsequent working
M0, M1	Method mark awarded 0, 1
A0, A1	Accuracy mark awarded 0, 1
B0, B1	Independent mark awarded 0, 1
E	Explanation mark 1
SC	Special case
^	Omission sign
MR	Misread
BP	Blank page
Highlighting	
Other abbreviations in	Meaning
mark scheme	
E1	Mark for explaining a result or establishing a given result
dep*	Mark dependent on a previous mark, indicated by *. The * may be omitted if only previous M mark.
cao	Correct answer only
oe	Or equivalent
rot	Rounded or truncated
soi	Seen or implied
www	Without wrong working
AG	Answer given
awrt	Anything which rounds to
BC	By Calculator
DR	This indicates that the instruction In this question you must show detailed reasoning appears in the question.

2. Subject-specific Marking Instructions for AS/A Level Further Mathematics B (MEI)

a Annotations must be used during your marking. For a response awarded zero (or full) marks a single appropriate annotation (cross, tick, M0 or ^) is sufficient, but not required.

For responses that are not awarded either 0 or full marks, you must make it clear how you have arrived at the mark you have awarded and all responses must have enough annotation for a reviewer to decide if the mark awarded is correct without having to mark it independently.

It is vital that you annotate standardisation scripts fully to show how the marks have been awarded.

Award NR (No Response)

- if there is nothing written at all in the answer space and no attempt elsewhere in the script
- OR if there is a comment which does not in any way relate to the question (e.g. 'can't do', 'don't know')
- OR if there is a mark (e.g. a dash, a question mark, a picture) which isn't an attempt at the question.

Note: Award 0 marks only for an attempt that earns no credit (including copying out the question).

If a candidate uses the answer space for one question to answer another, for example using the space for 8(b) to answer 8(a), then give benefit of doubt unless it is ambiguous for which part it is intended.

b An element of professional judgement is required in the marking of any written paper. Remember that the mark scheme is designed to assist in marking incorrect solutions. Correct solutions leading to correct answers are awarded full marks but work must not always be judged on the answer alone, and answers that are given in the question, especially, must be validly obtained; key steps in the working must always be looked at and anything unfamiliar must be investigated thoroughly. Correct but unfamiliar or unexpected methods are often signalled by a correct result following an apparently incorrect method. Such work must be carefully assessed. When a candidate adopts a method which does not correspond to the mark scheme, escalate the question to your Team Leader who will decide on a course of action with the Principal Examiner.

If you are in any doubt whatsoever you should contact your Team Leader.

c The following types of marks are available.

Μ

A suitable method has been selected and *applied* in a manner which shows that the method is essentially understood. Method marks are not usually lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. In some cases the nature of the errors allowed for the award of an M mark may be specified.

A method mark may usually be implied by a correct answer unless the question includes the DR statement, the command words "Determine" or "Show that", or some other indication that the method must be given explicitly.

Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated Method mark is earned (or implied). Therefore M0 A1 cannot ever be awarded.

В

Mark for a correct result or statement independent of Method marks.

Е

A given result is to be established or a result has to be explained. This usually requires more working or explanation than the establishment of an unknown result.

Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored. Sometimes this is reinforced in the mark scheme by the abbreviation isw. However, this would not apply to a case where a candidate passes through the correct answer as part of a wrong argument.

- d When a part of a question has two or more 'method' steps, the M marks are in principle independent unless the scheme specifically says otherwise; and similarly where there are several B marks allocated. (The notation 'dep*' is used to indicate that a particular mark is dependent on an earlier, asterisked, mark in the scheme.) Of course, in practice it may happen that when a candidate has once gone wrong in a part of a question, the work from there on is worthless so that no more marks can sensibly be given. On the other hand, when two or more steps are successfully run together by the candidate, the earlier marks are implied and full credit must be given.
- e The abbreviation FT implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A and B marks are given for correct work only differences in notation are of course permitted. A (accuracy) marks are not given for answers obtained from incorrect working. When A or B marks are awarded for work at an intermediate stage of a solution, there may be various alternatives that are equally acceptable. In such cases, what is acceptable will be detailed in the mark scheme. If this is not the case, please escalate the question to your Team Leader who will decide on a course of action with the Principal Examiner.

Sometimes the answer to one part of a question is used in a later part of the same question. In this case, A marks will often be 'follow through'. In such cases you must ensure that you refer back to the answer of the previous part question even if this is not shown within the image zone. You may find it

Mark Scheme

easier to mark follow through questions candidate-by-candidate rather than question-by-question.

f Unless units are specifically requested, there is no penalty for wrong or missing units as long as the answer is numerically correct and expressed either in SI or in the units of the question. (e.g. lengths will be assumed to be in metres unless in a particular question all the lengths are in km, when this would be assumed to be the unspecified unit.)

We are usually quite flexible about the accuracy to which the final answer is expressed; over-specification is usually only penalised where the scheme explicitly says so.

- When a value is given in the paper only accept an answer correct to at least as many significant figures as the given value.
- When a value is **not given** in the paper accept any answer that agrees with the correct value to **2 s.f.** unless a different level of accuracy has been asked for in the question, or the mark scheme specifies an acceptable range.

NB for Specification A the rubric specifies 3 s.f. as standard, so this statement reads "3 s.f"

Follow through should be used so that only one mark in any question is lost for each distinct accuracy error.

Candidates using a value of 9.80, 9.81 or 10 for g should usually be penalised for any final accuracy marks which do not agree to the value found with 9.8 which is given in the rubric.

- g Rules for replaced work and multiple attempts:
 - If one attempt is clearly indicated as the one to mark, or only one is left uncrossed out, then mark that attempt and ignore the others.
 - If more than one attempt is left not crossed out, then mark the last attempt unless it only repeats part of the first attempt or is substantially less complete.
 - if a candidate crosses out all of their attempts, the assessor should attempt to mark the crossed out answer(s) as above and award marks appropriately.
- h For a genuine misreading (of numbers or symbols) which is such that the object and the difficulty of the question remain unaltered, mark according to the scheme but following through from the candidate's data. A penalty is then applied; 1 mark is generally appropriate, though this may differ for some units. This is achieved by withholding one A or B mark in the question. Marks designated as cao may be awarded as long as there are no other errors. If a candidate corrects the misread in a later part, do not continue to follow through. E marks are lost unless, by chance, the given results are established by equivalent working. Note that a miscopy of the candidate's own working is not a misread but an accuracy error.
- i If a calculator is used, some answers may be obtained with little or no working visible. Allow full marks for correct answers provided that there is nothing in the wording of the question specifying that analytical methods are required such as the bold "In this question you must show detailed reasoning", or the command words "Show" and "Determine. Where an answer is wrong but there is some evidence of method, allow appropriate method marks. Wrong answers with no supporting method score zero. If in doubt, consult your Team Leader.
- j If in any case the scheme operates with considerable unfairness consult your Team Leader.

C	Juesti	on	Answer	Marks	AO	Guidance
1	(a)		1.15 – 0.85 or 1.05 – 0.95	M1	1.1a	both values attempted, one must be correct;
			$0.1 < x_2 - x_1 < 0.3$ oe isw	A1	1.1	allow non-strict inequality or eg 0.1 to 0.3 oe allow SC1 for correct answer unsupported
				[2]		
1	(b)		2.8 - 2.3 or 2.7 - 2.4	M1	1.1a	both values attempted, one must be correct
			$0.3 < y_2 - y_1 < 0.5$ oe isw	A1	1.1	allow non-strict inequality or eg 0.3 to 0.5 oe allow SC1 for correct answer unsupported
				[2]		
1	(c)		their $\frac{0.5}{0.1}$ and their $\frac{0.3}{0.3}$	M1	3.1 a	
			1 < m < 5 cao isw	A1	3.2a	
				[2]		
1	(d)		the denominator involves the subtraction	B1	2.4	must refer to denominator or to division by difference
			of nearly equal quantities oe			between two nearly equal numbers
				[1]		
2	(a)		because the values of <i>t</i> are not evenly spaced	B1	1.2	do not allow
			oe			eg because the data are not evenly spaced
						eg because the x values are not evenly spaced
				[1]		

(Questio	n Answer	Marks	AO	Guidance
2	(b)	$\frac{(t-0)(t-3)}{(5-0)(5-3)} \times 27.0 + \frac{(t-0)(t-5)}{(3-0)(3-5)} \times 29.4 +$	M1	3.3	allow sign errors in substitution of <i>x</i> -values and one numerical error
		$\frac{(t-3)(t-5)}{(0-3)(0-5)} \times 34.5$	A1	1.1	all substitutions correct
		$0.1t^2 - 2t + 34.5$ $P_2(t) = 0.1t^2 - 2t + 34.5$ or	A1	1.1	two terms correct
		$P = 0.1t^2 - 2t + 34.5$	A1	1.1	all correct; A0 if other variable used or <i>P</i> omitted; do not allow $f(t)$
			[4]		
2	(c)	$t = 6$ gives $P = 26.1 \approx 26.0$ so good fit	B1	3.4	if B0B0 allow SC1 for 26.1 and 24.5 seen
		$t = 10$ gives $P = 24.5 \approx 24.4$ so good fit	B1	3.4	may see one comment for both comparisons
			[2]		
2	(d)	model predicts tyre pressures will start to	B1	3.5b	allow
		increase (after $t = 10$)			eg as $t \to \infty, P \to \infty$
			[1]		
3	(a)	$\frac{2.318559 - 2.094395}{2.5 - 2} \text{ oe soi}$	M1	1.1	
		0.448328 isw or 0.44833 isw	A1	1.1	allow B2 for correct answer unsupported
		2 210550-1 602127	[2]		
3	(b)	$\frac{2.318559 - 1.682137}{2.5 - 1.5} $ oe soi	M1	1.1	
		0.636422 isw or 0.63642 isw	A1	1.1	allow B2 for correct answer unsupported
			[2]		

	Question	Answer	Marks	AO	Guidance
3	(c)	 chord joining point on curve where x = 1.5 to point on curve where x = 2.5 and tangent to the curve at x = 2 drawn; [and gradient of chord identified as an approximation to the gradient of the tangent] 	B1	1.1	
			[1]		
3	(d)	 central difference method is (usually) a second order method whereas forward difference is (usually) a first order method, so answer to part (b) probably more accurate <i>or</i> central difference method uses <i>x</i>-values on both sides of 2 oe whereas forward difference method uses a step in the positive 	B1	2.2b	must mention both methods must mention both methods
		<i>x</i> -direction only oe			
			[1]		
4	(a)	$x^{\frac{1}{x}}$ seen correct limits identified in integral	B1 B1	1.1 1.1	not necessarily as an integral
		$\int_{1.25}^{1.75} x^{\frac{1}{x}} \mathrm{d}x$	B1	1.1	all correct
					if B1B0B0 allow B1 SCB1 for $\int_a^b x^{\frac{1}{x}} dx$, where <i>a</i> and <i>b</i> are numerical values symmetrical about 1.5
			[3]		

(Questi	on	Answer	Marks	AO	Guidance
4	(b)		the curve is concave down so the midpoint rule will give an over-estimate	B1	2.4	or the curve is increasing and the gradient of the curve is decreasing so the midpoint rule will give an over-estimate
				[1]		
5	(a)		the gradient of $g(x)$ at the root is less than -1 oe so the iteration will not converge	B1	2.4	must refer to gradient at root or gradient of $g(x)$ at intersection of $[y =] x$ and $[y =] g(x)$; allow eg won't converge because $ g'(1.8) > 1$
				[1]		
5	(b)		2 1.81651017847 1.81159652649	B1 B1	1.1 2.1	1.816510 and 1.811596 oe seen 1.8115900 and 1.8115900 oe seen
			1.81159008685 1.81159008298 1.811590	B1	2.2a	dependent on award of first two marks if B1B0B0 allow B1 SCB2 for 1.811590 seen twice and 6 dp accuracy confirmed by change of sign
				[3]		
5	(c)		ratio of differences appears to converge to a constant [0.0006]	B1	2.2a	ignore comments on speed of convergence
			(which suggests) first order convergence	B1	2.2b	B0 if spoiled by incorrect reasoning
				[2]		
6	(a)	(i)	$\frac{(3.14^2+2)-(\pi^2+2)}{(\pi^2+2)} \text{ or } \frac{(3.14^2+2)}{(\pi^2+2)} - 10e$	M1	1.1	allow omission of 2 in numerator; allow bracket error; may be embedded in modulus
			awrt -0.000 843 cao	A1	1.1	NB -8.43×10^{-4} allow B2 for correct answer unsupported

Question		on	Answer	Marks	AO	Guidance
				[2]		
6	(a)	(ii)	it would be different as the order of operations is different	B1	2.4	
				[1]		
6	(b)	(i)	1.00001×10^{-12}	B1	1.1	or 0.000 000 000 001 000 01
				[1]		
6	(b)	(ii)	not zero because the value stored in B2 and B3 is not stored as 1, even though they are displayed as 1	B1	2.4	do not allow just eg the spreadsheet stores values to greater accuracy than it displays
			not equal to the [values in A2 and A3] because the values in B2 and B3 are different to each other (even though they are displayed as 1), so different values are used to compute the values in C2 and C3	B1	2.4	do not allow just eg the spreadsheet stores values to greater accuracy than it displays
				[2]		
7	(a)		use of $f(2 + 0.03) \approx f(2) + 0.03f'(2)$ soi	M1	1.2	may see eg $\pm 2.8 \times 0.03$ may be implied by 5.916
			(error ≈) –0.084 cao	A1	1.1	mark the final answer allow SC1 for correct final answer unsupported
				[2]		

Q	Juesti	on	Answer	Marks	AO	Guidance
7	(b)	(i)	ratio of differences is decreasing	B1	2.2a	ignore superfluous comments
			(which suggests) convergence is faster than first order; allow (which suggests) convergence is higher than first order	B1	2.2b	 do not allow if spoiled by incorrect reasoning; do not allow (which suggests) second order convergence do not allow greater than first order convergence if B0B0 allow SC1 for ratio of differences not converging to a constant so convergence not first order
				[2]		
7	(b)	(ii)	2.182419066 since convergence is faster than first order / generally second order	B1 B1	2.2b 2.4	allow 2.18241907 or 2.1824191 or allow B1 for 2.182419 and B1 for eg since x_4 and x_5 agree to this precision oe; allow last two estimates or values in B6 and B7
				[2]		
8	(a)			B1	1.1	chord from curve at $x = 1$ and through the curve at $x = 2$ which joins the <i>x</i> -axis to the right of the root isw
				[1]		
8	(b)		= L3	B1	1.1	
				[1]		

Question		Answer	Marks	AO	Guidance	
8	(c)	= (J3*M3 - K3*L3)/(M3 - K3)	M1	1.1	use of secant formula; allow spreadsheet syntax errors but not error in formula NB $rr (12*K4 - K2*L2)/(K4 - K2)$	
		or = $(K3*L3 - J3*M3)/(K3 - M3)$ oe			NB or = $(J3*K4 - K3*L3)/(K4 - K3)$ or = $(K3*L3 - J3*K4)/(K3 - K4)$	
					or (J3*K4 – K3*J4)/(K4 – K3)	
					or = $(K3*J4 - J3*K4)/(K3 - K4)$	
			A1	1.1	all correct	
			[2]			
8	(d)	2.12664	B1	2.2a		
			[1]			
8	(e)	f(2.126635) = awrt - 0.00007	M1	2.1	both calculated and one correct to 1 sf	
					NB –0.000070377215 and –0.000066109812	
					M1FT 2.13042	
					awrt 0.002 in each case	
					NB 0.0015487 and 0.001553	
		f(2.126645) = awrt - 0.00007				
		no sign change so not correct to 5 dp	A1	2.2a	both values correct to 1 sf	
					A1 FT awrt 0.002 and 0.002 so no sign change so not	
					correct to 5 dp	
			[2]			

(Questio	on Answer	Marks	AO	Guidance
8	(f)	$1 < a \le 1.74$	B 1	3.1a	
			[1]		
9	(a)	$\frac{4 \times 0.66617652 - 0.52764369}{3} \mathbf{0e} \text{ or}$	M1	1.1	
		$\frac{4 \times 0.72534275 - 0.66617652}{3}$ oe			
		$[S_2 =] 0.712354$ to 0.71235413 and	A1	1.1	allow A1 if not attributed, but A0 if wrongly attributed
		[S ₄ =] 0.7450648 to 0.745065			NB 0.745064826667
					allow SC1 for both answers correct unsupported
			[2]		
9	(b)	0.7 because the two Simpson's estimates	B1	2.2b	or 0.8 because the approximations are increasing and the
		only agree to this precision			difference between S_4 and S_2 is large enough to suggest value closer to 0.8 than 0.7
			[1]		
9	(c)	ratio of differences seems to be converging [to 0.4]	B1	2.2b	
		0.25 < r < 0.5 hence order of (convergence of) method is between first and second order (in this case)	B1	2.2b	Do not allow eg order of convergence is between 1 and 2
			[2]		

Mark Scheme

(Questi	on	Answer	Marks	AO	Guidance
9	(d)		$0.7655404 + 0.001618 \times \frac{0.4}{1 - 0.4}$	M1	3.1 a	extrapolation with values of T_n , associated difference and r deduced from table; allow for partial extrapolation
				A1	1.1	allow 0.4 to 0.401 for <i>r</i> ; allow for partial extrapolation NB 0.7661876 with <i>r</i> = 0.4 NB 0.7661891371 with <i>r</i> = 0.40095
			0.766619 to 0.766624	A1	1.1	NB 0.766619066667 and 0.76662334316 from extrapolation to infinity with 0.4 and 0.40095 respectively
			0.767 or 0.7666 since extrapolation much more accurate oe	A1	3.2a	allow 0.77 is secure by comparison of T_{64} with extrapolated value oe
						if M0 , allow SC2 for extrapolation with $r = 0.25$ to obtain 0.766079
				[4]		

APPENDIX

Exemplar responses for Q2(b)

Response	Mark

Need to get in touch?

If you ever have any questions about OCR qualifications or services (including administration, logistics and teaching) please feel free to get in touch with our customer support centre.

Call us on

01223 553998

Alternatively, you can email us on

support@ocr.org.uk

For more information visit





Twitter/ocrexams

/ocrexams

/company/ocr

/ocrexams



OCR is part of Cambridge University Press & Assessment, a department of the University of Cambridge.

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored. © OCR 2023 Oxford Cambridge and RSA Examinations is a Company Limited by Guarantee. Registered in England. Registered office The Triangle Building, Shaftesbury Road, Cambridge, CB2 8EA.

Registered company number 3484466. OCR is an exempt charity.

OCR operates academic and vocational qualifications regulated by Ofqual, Qualifications Wales and CCEA as listed in their qualifications registers including A Levels, GCSEs, Cambridge Technicals and Cambridge Nationals.

OCR provides resources to help you deliver our qualifications. These resources do not represent any particular teaching method we expect you to use. We update our resources regularly and aim to make sure content is accurate but please check the OCR website so that you have the most up-to-date version. OCR cannot be held responsible for any errors or omissions in these resources.

Though we make every effort to check our resources, there may be contradictions between published support and the specification, so it is important that you always use information in the latest specification. We indicate any specification changes within the document itself, change the version number and provide a summary of the changes. If you do notice a discrepancy between the specification and a resource, please <u>contact us</u>.

Whether you already offer OCR qualifications, are new to OCR or are thinking about switching, you can request more information using our Expression of Interest form.

Please get in touch if you want to discuss the accessibility of resources we offer to support you in delivering our qualifications.